arXiv:1612.08158v1 [math.NT] 24 Dec 2016

$(1^3+1)(2^3+1)\cdots(n^3+1)$ is not a cube*

Chuanze Niu†

Abstract

For a positive integer n, define

$$C_n = \prod_{k=1}^{n}(k^3+1).$$

In this paper we prove that there are no cubes in the integer sequence $C_n,\ n = 1, 2, \cdots$.

Keywords: Cubes, the p-adic orders, the distribution of prime numbers
MSC2003: 11D25; 11D45

1 Introduction

Let $f(x) = ax^d + b$ be a polynomial of degree d in $\mathbb{Z}[x]$. Define the consecutive product of the sequence $f(k),\ k = 1, 2, \cdots$, by $C_n(f) = \prod_{k=1}^{n} f(k)$. It is interesting to discuss whether $C_n(f)$ is a d-th power of integers or not.

When $f(x) = x^2 + 1$, Javier Cilleruelo [1] proved that $C_n(f)$ is a square only for $n = 3$ using the analytic methods and answered the conjecture proposed by [2]. More generally, Yang, et al. [3] showed that there are finitely many squares in $C_n(f)$ when $d = 2$ and $(a, b) = 1$, $1 \le a \le 10$ and $1 \le b \le 20$. Under ABC conjecture, Zhang [4] proved $C_n(f)$ is not a d-th power for n large enough, where $d = 2^l \cdot 3^m$ with $l \ge 1, l + m \ge 2, ab \ne 0$ and such that $f(k) \ne 0$ for any $k \ge 1$. When $f(x) = x^2 - 1$, Hong [5] proved that there are infinitely many n such that $C_n(f)$ is a square using Dirichlet's unit theorem.

As can be seen, when f is of degree 2, there exists many celebrate results. In this paper, we assume $f(x) = x^3 + 1$ and let $C_n = C_n(f)$, drawing on the idea of Cilleruelo [1], we show that C_n is never a cube, the main result is the following theorem.

Theorem 1.1. *For any positive integer n, the value $C_n = (1^3+1)(2^3+1)\cdots(n^3+1)$ is not a cube.*

As a corollary of the main theorem, we get the following result.

Corollary 1.2. *The Diophantine equation*

$$y^3 = \prod_{k=1}^{n}(k^3 + a^3)$$

has no solutions with $a = 1$.

*Supported by National Natural Science Foundation of China (Grant No. 11401285)

†School of Mathematical Sciences, Liaocheng University, Liaocheng 252059, P. R. China, E-mail: niuchuanze@lcu.edu.cn

2 Preliminaries

In this section, we do some preparations that used to prove the main theorem of this paper which include the upper bound of the p-adic order of C_n and the upper bound of the distribution function $\pi(x)$ of primes. We always assume $n \geq 4$ since one may check $C_1,\ C_2$ and C_3 are not cubes easily.

Let p be a rational prime number, the p-adic order of a positive integer n is denoted by $\mathrm{ord}_p(n)$, then $n = \prod p^{\mathrm{ord}_p(n)}$, where the product runs through all prime factors p of n. Note that

$$\mathrm{ord}_p(n) = \sum_{1 \leq j \leq \frac{\log n}{\log p}} \chi(p^j | n),$$

where $\chi(p^j|n)$ equals 1 or 0 depending on whether n is divisible by p^j or not, then

$$ord_p(C_n) = \sum_{1 \leq j \leq \frac{\log(n^3+1)}{\log p}} \sharp\{1 \leq k \leq n |\ p^j | (k^3+1)\}. \tag{2.1}$$

Lemma 2.1. *If $p = n+1$ is a prime factor of C_n, then $ord_p(C_n) = 1$.*

Proof. Since $p = n+1$, we have $ord_p(n^3+1) \geq 1$. If $ord_p(n^3+1) > 1$ then $p|(n^2-n+1)$ and therefore p divides $(n+1)^2 - (n^2-n+1) = 3n$, which is a contradiction since $p > n > 3$. We claim that there are no integers k with $1 \leq k < n$ such that $p|((n-k)^3+1)$. Otherwise, p divides $n^3+1-(n-k)^3-1 = 3nk(n-k)$, which is impossible. Therefore,

$$ord_p(C_n) = ord_p(n^3+1) = 1.$$

This completes the proof. □

In the proof of Lemma 2.1, we see in the sequence $k^3+1,\ k = 1, 2, \cdots$, the first term that divisible by a prime p is $(p-1)^3+1$.

Lemma 2.2. *If $p > n+1$ is a prime factor of C_n, then $ord_p(C_n) \leq 2$.*

Proof. Note that $p > k+1$ and $p^2 > k^2-k+1$, we see p^2 can not divide k^3+1 for positive integers $k \leq n$. Therefore $\mathrm{ord}_p(k^3+1) = 0$ or 1.

If $p|(k^3+1)$ and $p|(l^3+1)$ for $1 \leq l < k \leq n$, then $p|(k^2-k+1)$ and $p|(l^2-l+1)$ which yields that $p|(k-l)(k+l-1)$. Therefore $p|(k+l-1)$ and it leads to $k+l-1 = p$ since $1 \leq k+l-1 < 2p$. So k and l are determined by each other. The number of those k such that $p|(k^3+1)$ is at most 2, thus by Equation (2.1), $\mathrm{ord}_p(C_n) \leq 2$. The proof is done. □

Proposition 2.3. *Let p be a prime factor of a cube C_n, then $p \leq n$.*

Proof. Note that for a prime factor p of a cube C_n, $\mathrm{ord}_p(C_n) \geq 3$. The proposition follows from Lemmas 2.1 and 2.2. □

By Proposition 2.3, one can rewrite a cube C_n as the product of primes $p \leq n$, i.e.,

$$C_n = \prod_{p \leq n} p^{\mathrm{ord}_p(C_n)}.$$

Next we always assume $f(x) = x^3+1$ and we shall discuss the number of the roots of $f(x) \equiv 0 (\mathrm{mod}\ p^j)$ for any positive integer j.

Lemma 2.4. *$f(x)(mod\ p)$ has 3 different roots if $p \equiv 1(mod\ 3)$ while it has exactly 1 root if $p \equiv 2(mod\ 3)$.*

Proof. Note that $x \equiv -1(\text{mod } p)$ is a trivial solution of $f(x) \equiv 0(\text{mod } p)$ and it is the unique solution if $p = 2$. For $p \geq 5$, consider the congruence

$$x^2 - x + 1 \equiv 0(\text{mod } p). \tag{2.2}$$

It is equivalent to

$$(2x - 1)^2 \equiv -3(\text{mod } p). \tag{2.3}$$

Therefore Equation (2.2) has no solutions if the Lengerde symbol $\left(\frac{-3}{p}\right) = -1$, while it has 2 different solutions if $\left(\frac{-3}{p}\right) = 1$. By Gauss reciprocity law,

$$\left(\frac{-3}{p}\right) = \left(\frac{-1}{p}\right)\left(\frac{3}{p}\right) = (-1)^{\frac{p-1}{2}}(-1)^{\frac{p-1}{2}}\left(\frac{p}{3}\right) = \left(\frac{p}{3}\right).$$

The lemma follows since $\left(\frac{p}{3}\right)$ equals -1 if $p \equiv 2(\text{mod } 3)$ and equals 1 if $p \equiv 1(\text{mod } 3)$. □

Lemma 2.5. *If $p \neq 3$, then the number of the roots of $f(x)(mod\ p)$ equals to the number of the roots of $f(x)(mod\ p^j)$ for any positive integer j.*

Proof. Note that $f(x) = x^3 + 1 \equiv 0(\text{mod } p)$ and $f'(x) = 3x^2 \equiv 0(\text{mod } p)$ have no common solutions if $p \neq 3$, then the lemma follows by Hensel' Lemma (or see [6], section 8.3). □

Lemma 2.6. *$x^3 + 1 \equiv 0(mod\ 3^j)$ has 3 different roots for any integer $j \geq 2$.*

Proof. Work by induction on j.

When $j = 2$, the roots of $x^3 + 1 \equiv 0(\text{mod } 3^2)$ are $x \equiv 2,\ 5,\ 8(\text{mod } 3^2)$.

Assume the roots of $x^3 + 1 \equiv 0(\text{mod } 3^j)$ are $x \equiv 3^{j-1} - 1,\ 2 \cdot 3^{j-1} - 1,\ 3^j - 1(\text{mod } 3^j)$. Next we show the roots of $x^3 + 1 \equiv 0(\text{mod } 3^{j+1})$ are $x \equiv 3^j - 1,\ 2 \cdot 3^j - 1,\ 3^{j+1} - 1(\text{mod } 3^{j+1})$. It suffices to show that $\xi^3 + 1 \not\equiv 0(\text{mod } 3^{j+1})$ when $\xi = 3^{j-1} - 1,\ 2 \cdot 3^{j-1} - 1$. Take $\xi = 3^{j-1} - 1$ as an example and the other case could be checked similarly. Note that $3j - 3 \geq j + 1$ and $2j - 1 \geq j + 1$,

$$(3^{j-1} - 1)^3 + 1 = 3^{3j-3} - 3^{2j-1} + 3^j \equiv 3^j \not\equiv 0(\text{mod } 3^{j+1}).$$

This completes the proof of Lemma 2.6. □

Remark 1. *In fact, we can also show for any prime p and positive integer j, $x^p + 1 \equiv 0(mod\ p^{j+1})$ has exactly p different roots $x \equiv ip^j - 1(mod\ p^{j+1}),\ 1 \leq i \leq p$.*

Proposition 2.7. *Let i be any integer. For any positive integer j, $f(x) \equiv 0(mod\ p^j)$ has only 1 solution in the interval $[i, i + p^j - 1]$ if $p \equiv 2(mod\ 3)$, whereas it has at most 3 different solutions in $[i, i + p^j - 1]$ for other p's.*

Proof. The proposition follows from Lemmas 2.4, 2.5 and 2.6. □

Remark 2. *In the sequence $k^3 + 1,\ k = 1,\ 2,\ \cdots$, when $p \equiv 2(mod\ 3)$, the first three terms that divisible by p are $(p-1)^3 + 1$, $(2p-1)^3 + 1$ and $(3p-1)^3 + 1$. Therefore C_n is not a cube if $p - 1 \leq n \leq 3p - 2$.*

As an immediate consequence of Equation (2.1) and Proposition 2.7, we get the upper bound of $\mathrm{ord}_p(C_n)$.

Proposition 2.8. *We have*

$$ord_p(C_n) \leq \begin{cases} \sum\limits_{1\leq j\leq \frac{\log(n^3+1)}{\log p}} \left\lceil \frac{n}{p^j} \right\rceil, & p \equiv 2(mod\ 3); \\ 3 \sum\limits_{1\leq j\leq \frac{\log(n^3+1)}{\log p}} \left\lceil \frac{n}{p^j} \right\rceil, & p \not\equiv 2(mod\ 3), \end{cases}$$

where $\lceil x \rceil$ denotes the least integer that is greater than or equals to x.

The inequality $C_n > (n!)^3$ which is the technique that is used in [1], that is, combining Proposition 2.8, we can deduce a bound for the sum of $\frac{\log p}{p-1}$ for prime numbers $p \leq n$ in arithmetical progression when C_n is a cube. We end this section by recalling a result on the distribution of prime numbers. Let $\pi(x)$ be the number of primes which do not exceed x.

Lemma 2.9. *If $x > 10000$, then*

$$\pi(x) < \frac{1.139x}{\log x}.$$

Proof. See Lemma 2.5 in [3]. □

3 Proof of Theorem 1.1

It is easy to see that

$$\mathrm{ord}_p(n!) = \sum_{1\leq j\leq \frac{\log n}{\log p}} \left[\frac{n}{p^j} \right],$$

where $[x]$ is the greatest integer not exceeding x. When $p \leq n$, we have

$$\mathrm{ord}_p(n!) \geq \sum_{1\leq j\leq \frac{\log n}{\log p}} \left(\frac{n}{p^j} - 1 \right) \leq \frac{n-p}{p-1} - \frac{\log n}{\log p} \leq \frac{n-1}{p-1} - \frac{2\log n}{\log p}. \tag{3.1}$$

Proof of Theorem 1.1. Note that

$$\prod_{p\leq n} p^{\mathrm{ord}_p(C_n)} > \prod_{p\leq n} p^{3\mathrm{ord}_p(n!)},$$

take logarithm on both sides, we get

$$\sum_{p\leq n} \mathrm{ord}_p(C_n) \log p > 3 \sum_{p\leq n} \mathrm{ord}_p(n!) \log p.$$

Then

$$\sum_{\substack{p\leq n \\ p\equiv 2(\mathrm{mod}\ 3)}} (\mathrm{ord}_p(n!) - \frac{1}{3}\mathrm{ord}_p(C_n)) \log p < \sum_{\substack{p\leq n \\ p\not\equiv 2(\mathrm{mod}\ 3)}} (\frac{1}{3}\mathrm{ord}_p(C_n) - \mathrm{ord}_p(n!)) \log p.$$

By Proposition 2.8 and Equation (3.1), when $p \not\equiv 2 (\mathrm{mod}\ 3)$, we have

$$\begin{aligned}
\frac{1}{3}\mathrm{ord}_p(C_n) - \mathrm{ord}_p(n!) &\leq \sum_{1\leq j\leq \frac{\log(n^3+1)}{\log p}} \left\lceil \frac{n}{p^j} \right\rceil - \sum_{1\leq j\leq \frac{\log n}{\log p}} \left\lfloor \frac{n}{p^j} \right\rfloor \\
&\leq \sum_{1\leq j\leq \frac{\log n}{\log p}} \left(\left\lceil \frac{n}{p^j} \right\rceil - \left\lfloor \frac{n}{p^j} \right\rfloor \right) + \sum_{\frac{\log n}{\log p} < j\leq \frac{\log(n^3+1)}{\log p}} \left\lceil \frac{n}{p^j} \right\rceil \\
&\leq \sum_{1\leq j\leq \frac{\log n}{\log p}} 1 + \sum_{\frac{\log n}{\log p} < j\leq \frac{\log(n^3+1)}{\log p}} 1 \\
&\leq \frac{\log(n^3+1)}{\log p};
\end{aligned}$$

when $p \equiv 2 (\mathrm{mod}\ 3)$, we have

$$\begin{aligned}
\mathrm{ord}_p(n!) - \frac{1}{3}\mathrm{ord}_p(C_n) &\geq \sum_{1\leq j\leq \frac{\log n}{\log p}} \left\lfloor \frac{n}{p^j} \right\rfloor - \frac{1}{3} \sum_{1\leq j\leq \frac{\log(n^3+1)}{\log p}} \left\lceil \frac{n}{p^j} \right\rceil \\
&= \frac{2}{3} \sum_{1\leq j\leq \frac{\log n}{\log p}} \left\lfloor \frac{n}{p^j} \right\rfloor + \frac{1}{3} \sum_{1\leq j\leq \frac{\log n}{\log p}} (\left\lfloor \frac{n}{p^j} \right\rfloor - \left\lceil \frac{n}{p^j} \right\rceil) - \frac{1}{3} \sum_{\frac{\log n}{\log p} < j\leq \frac{\log(n^3+1)}{\log p}} \left\lceil \frac{n}{p^j} \right\rceil \\
&\geq \frac{2}{3} \cdot \frac{n-1}{p-1} - \frac{4}{3} \cdot \frac{\log n}{\log p} - \frac{1}{3} \cdot \frac{\log(n^3+1)}{\log p}.
\end{aligned}$$

Therefore, we have

$$\begin{aligned}
\frac{2(n-1)}{3} \sum_{\substack{p\leq n \\ p\equiv 2(\mathrm{mod}\ 3)}} \frac{\log p}{p-1} &\leq \sum_{\substack{p\leq n \\ p\not\equiv 2(\mathrm{mod}\ 3)}} \log(n^3+1) + \frac{4}{3} \sum_{\substack{p\leq n \\ p\equiv 2(\mathrm{mod}\ 3)}} \log n + \frac{1}{3} \sum_{\substack{p\leq n \\ p\equiv 2(\mathrm{mod}\ 3)}} \log(n^3+1) \\
&\leq \frac{2}{3} \sum_{\substack{p\leq n \\ p\not\equiv 2(\mathrm{mod}\ 3)}} \log(n^3+1) + \frac{4}{9} \sum_{\substack{p\leq n \\ p\equiv 2(\mathrm{mod}\ 3)}} \log(n^3+1) + \frac{1}{3} \sum_{p\leq n} \log(n^3+1) \\
&= \frac{2}{9} \sum_{\substack{p\leq n \\ p\not\equiv 2(\mathrm{mod}\ 3)}} \log(n^3+1) + \frac{4}{9} \sum_{p\leq n} \log(n^3+1) + \frac{1}{3} \sum_{p\leq n} \log(n^3+1) \\
&\leq \pi(n) \log(n^3+1).
\end{aligned}$$

Therefore

$$\sum_{\substack{p\leq n \\ p\equiv 2(\mathrm{mod}\ 3)}} \frac{\log p}{p-1} \leq \frac{3}{2(n-1)} \pi(n) \log(n^3+1).$$

Now using the upper bound of $\pi(x)$ in Lemma 2.9, when $n > 10000$,

$$\sum_{\substack{p\leq n \\ p\equiv 2(\mathrm{mod}\ 3)}} \frac{\log p}{p-1} \leq \frac{3.417}{2} \cdot \frac{n}{n-1} \cdot \frac{\log(n^3+1)}{\log n}.$$

After computations, the right-hand side is < 5.2 if $n \geq 10000$. Adding over enough prime numbers for $p \equiv 2(\text{mod } 3)$, we see that for $n \geq 34631$,

$$\sum_{\substack{p \leq n \\ p \equiv 2(\text{mod } 3)}} \frac{\log p}{p-1} \geq 5.2.$$

Hence C_n is not a cube when $n \geq 34631$. For $4 \leq n \leq 34631$, by Remark 2, we consider the following primes that congruent to 2 modulo 3,

$$5,\ 11,\ 29,\ 83,\ 233,\ 683,\ 2039,\ 6113,\ 18329,$$

which corresponding intervals such that C_n are not cubes if n lies in them:

$$[4, 13],\ [10, 31],\ [28, 85],\ [83, 247],\ [232, 697],\ [682, 2047],\ [2038, 6115],\ [6112, 18337],\ [18328, 54985].$$

Therefor C_n is not a cube if $4 \leq n \leq 54985$, this completes the proof.

References

[1] J.Cilleruelo, Squares in $(1^2+1)\cdots(n^2+1)$, J. Number Theory 128, 2008, 2488-2491.

[2] T. Amdeberhan, et al., Arithmetical propeties of a sequence arising from an arctangent sum, J. Number Theory 128(6), 2008, 1807-1846.

[3] Yang S, et al., Diophantine equations with products of consecutive values of a quadratic polynomial, J. Number Theory. 2011, 131: 1840-1851.

[4] Z.Zhang, Powers in $\prod_{k=1}^{n}(ak^{2^l \cdot 3^m}+b)$, Functiones Et Approximatio Commentarii Mathematici, 2012, 46(1): 7-13.

[5] Shaofang Hong and Xingjiang Liu, Squares in $(2^2-1)\cdots(n^2+1)$ and p-adic valuation, Asian-Europe J. Math, 2010, 3(2): 329.

[6] G. Hardy, E.Wright. An introducton to the theory of numbers, sixth edition, Oxford Univ. Press, 2009.